

60th Annual Scientific Session & Expo

E70

JACC April 5, 2011

Volume 57, Issue 14



CARDIAC ARRHYTHMIAS

TWELVE-LEAD ECG PATTERNS FAIL TO IDENTIFY AN EPICARDIAL ORIGIN FOR LEFT VENTRICULAR TACHYCARDIA IN POST-INFARCTION PATIENTS

ACC Poster Contributions

Ernest N. Morial Convention Center, Hall F

Sunday, April 03, 2011, 3:30 p.m.-4:45 p.m.

Session Title: Clinical Electrophysiology --Ventricular Arrhythmias

Abstract Category: 27. Clinical Electrophysiology--Ventricular Arrhythmias

Session-Poster Board Number: 1057-376

Authors: *Martin Martinek, Usha B. Tedrow, Keiichi Inawa, Michifumi Tokuda, William G. Stevenson, Brigham and Women's Hospital, Boston, MA, Harvard Medical School, Boston, MA*

BACKGROUND Several ECG features have been reported to identify epicardial origins for left ventricular tachycardias (LV-VTs) in the absence of myocardial infarction. Only limited data exist in post-infarction patients. We tested proposed algorithms for non-ischemic tachycardias for their ability to identify epicardial LV-VT origins.

METHODS The QRS features of 17 successful epicardial and 29 endocardial RBBB LV-VTs were retrospectively reviewed by four independent electrophysiologists and analyzed for various 12-lead ECG features.

RESULTS All 12-lead ECG features proposed for non-ischemic LV-VTs were unable to consistently predict an epicardial LV-VT origin in infarct-related tachycardias. QRS duration was 199 ± 31 ms in epicardial vs. 178 ± 40 ms in endocardial LV-VTs ($p=0.063$) showing significant overlap and therefore lacking a reasonable cutoff-value between epicardial and endocardial LV-VTs. Pseudo-delta duration was 47 ± 9 vs. 54 ± 21 ms ($p=ns$), intrinsicoid deflection time was 103 ± 32 vs. 92 ± 24 ms ($p=ns$), shortest RS was 110 ± 36 vs. 101 ± 34 ms ($p=ns$) and median deflection index was 0.75 ± 0.27 vs. 0.85 ± 0.25 ms ($p=ns$).

The finding of a Q wave in lead I and absence of a Q wave in the inferior leads failed to predict an epicardial origin in superior LV-VT sites. Q waves in any inferior lead as well as an aVR/aVL ratio < 1 were not specific for an epicardial origin in inferior sites (all $p=ns$). Furthermore, most inferior LV-VTs showed a Q wave in the inferior leads which correlated with pre-existing Q-waves in sinus rhythm ECGs ($p=0.045$) rendering this ECG feature not useful.

CONCLUSION Proposed 12-lead ECG algorithms for the differentiation of epicardial vs. endocardial sites for non-ischemic LV-VTs are not valid in post-infarction patients.